

## PROPORTION OF DEATHS FROM TYPHOID FEVER PER 1,000 OF POPULATION IN THE FOLLOWING CITIES IN 1888.

CITIES.	Per 1,000.	CITIES.	Per 1,000.
Woodstock.....	1.20	Ottawa.....	0.42
Hull.....	1.16	Victoria, B.C.....	0.40
Brantford.....	0.99	Three Rivers.....	0.40
Winnipeg.....	0.95	Sorel.....	0.38
St. Hyacinthe.....	0.82	Halifax.....	0.35
Galt.....	0.69	Hamilton.....	0.35
Sherbrooke.....	0.65	Quebec.....	0.33
Montreal.....	0.56	Toronto.....	0.30
Belleville.....	0.55	St. Thomas.....	0.29
Charlottetown.....	0.51	St. John.....	0.11

The proportion of deaths from typhoid fever in the aggregate population of the 27 cities making returns was 0.44 per 1,000 persons living as compared with 0.55 in 1887.

136. In the cities of Ontario in 1887 the proportion was 0.54 per 1,000, while for the whole province it was 2.68 per 10,000 persons living, which compares very favourably with some other British possessions as the following figures show :—

Deaths from typhoid fever in Ontario & certain countries.

## DEATHS FROM TYPHOID FEVER PER 10,000 INHABITANTS IN THE FOLLOWING COUNTRIES.

1887—England and Wales.....	1.82
1888—Ireland.....	3.00
1884—Scotland.....	2.70
1887—Ontario.....	2.68
1886—Victoria.....	5.30
1886—New South Wales.....	6.03
1886—Queensland.....	12.27
1886—South Australia.....	3.39
1886—Tasmania.....	3.47
1886—New Zealand.....	2.30

137. The proportion of deaths from zymotic or specially preventable diseases to the total number of deaths was 22.00 per cent. as compared with 23.71 per cent. in 1887. The proportion to the aggregate population was 0.48 per cent., or 4.88 per 1,000 persons living, and the following table shows how they were distributed among the various cities.

Deaths in Canada from zymotic diseases.