

## DEATHS FROM TYPHOID FEVER IN THE UNDERMENTIONED CITIES IN 1887, 1888 AND 1889.

CITIES.	1887.	1888.	1889.	CITIES.	1887.	1888.	1889.
Montreal.....	125	113	87	Peterboro'.....	6	1	.....
Toronto.....	71	51	49	Fredericton.....	1	1	1
Quebec.....	22	21	40	St. Hyacinthe.....	6	6	9
Hamilton.....	19	15	17	Galt.....	.....	5	.....
Halifax.....	6	14	11	Hull.....	20	14	4
Winnipeg.....	21	21	38	London.....	8	7	4
Ottawa.....	43	17	19	Brantford.....	9	13	12
St. John, N.B.....	6	3	11	Victoria, B.C.....	8	6	6
Kingston.....	1	6	7	Three Rivers.....	2	4	4
St. Thomas.....	1	3	3	Woodstock.....	8	10	9
Charlottetown.....	4	7	8	Sorel.....	3	3	4
Guelph.....	3	2	2	Windsor.....	.....	1	2
Belleville.....	4	6	1	St. Johns, Que.....	.....	.....	3
Chatham.....	8	2	6				
Sherbrooke.....	6	6	4	Total.....	411	358	361

## PROPORTION OF DEATHS FROM TYPHOID FEVER PER 1,000 OF POPULATION IN THE FOLLOWING CITIES IN 1889.

CITIES.	Per 1,000.	CITIES.	Per 1,000.
Winnipeg.....	1.48	Kingston.....	0.38
St. Hyacinthe.....	1.12	Three Rivers.....	0.38
Woodstock, Ont.....	1.02	Victoria, B.C.....	0.32
Brantford.....	0.90	Hull.....	0.28
Chatham, Ont.....	0.70	St. Thomas.....	0.28
Quebec.....	0.60	Toronto.....	0.27
Charlottetown.....	0.58	Halifax.....	0.26
St. Johns, Que.....	0.54	St. John, N.B.....	0.25
Sorel.....	0.48	Guelph.....	0.19
Ottawa.....	0.43	Windsor.....	0.19
Sherbrooke.....	0.42	London.....	0.15
Montreal.....	0.41	Fredericton.....	0.13
Hamilton.....	0.38	Belleville.....	0.08

It will be seen that the death rate is considerably higher in many of the cities than it ought to be, though there was a slight decrease in the proportion to the aggregate population of the 28 cities included in the table, this having been 0.41, as compared with 0.44 in 1888 and 0.55 in 1887. The proportion in Ontario in 1888 was 0.28 per 1,000 inhabitants. In England, in the same year, the rate was, for the whole country 0.17, and for London 0.16, showing what can be accomplished by care and attention to sanitary matters.