

DEATHS FROM TYPHOID FEVER IN THE UNDERMENTIONED
PLACES IN 1887, 1888, 1889, 1890 AND 1891.

Cities and Towns.	1887.	1888.	1889	1890	1891.	Cities and Towns.	1887	1888	1889	1890	1891
Montreal	125	113	87	76	74	Fredericton	1	1	1	3	—
Toronto	71	51	49	117	117	St. Hyacinthe	6	6	9	3	2
Quebec	22	21	40	20	19	Galt	5	5	1	2	2
Hamilton	19	15	17	23	10	Hull	20	14	4	4	2
Halifax	6	14	11	7	8	London	8	7	4	6	8
Winnipeg	21	21	38	28	19	Brantford	9	13	12	11	8
Ottawa	43	17	19	19	9	Victoria, B.C.	8	6	6	6	13
St. John, N.B.	6	3	11	10	10	Three Rivers	2	4	4	3	5
Kingston	1	6	7	10	11	Woodstock	8	10	9	3	4
St. Thomas	1	3	3	7	3	Sorel	3	3	4	1	1
Charlottetown	4	7	8	3	7	Windsor	1	2	3	4	4
Guelph	3	2	2	3	2	St. John's, Que	3	3	5	3	3
Belleville	4	6	1	3	2	Brockville	1	1	3	1	3
Chatham	8	2	6	4	5	Stratford	1	1	1	1	1
Sherbrooke	6	6	4	1	6	Total	411	358	361	382	363
Peterboro'	6	1	2	5						

PROPORTION OF DEATHS FROM TYPHOID FEVER PER 1,000 OF
POPULATION IN THE FOLLOWING PLACES IN 1891.

Cities and Towns.	Per 1,000.	Cities and Towns.	Per 1,000.
Victoria, B.C.	0.77	Quebec	0.30
Winnipeg	0.74	St. Thomas	0.28
Toronto	0.64	St. Hyacinthe	0.28
St. John's, Que	0.62	Galt	0.26
Charlottetown	0.61	St. John, N.B.	0.25
Sherbrooke	0.59	London	0.25
Three Rivers	0.59	Belleville	0.20
Kingston	0.57	Ottawa	0.20
Chatham	0.55	Hamilton	0.20
Brantford	0.52	Guelph	0.18
Peterboro'	0.51	Hull	0.17
Woodstock, Ont	0.46	Halifax	0.17
Windsor, Ont	0.38	Sorel	0.14
Brockville	0.34	Stratford	0.10
Montreal	0.34		

176. The following table shows how the assigned causes of death, in the places making returns, were distributed among the several classes. The classification is based upon that established by the Registrar-General of England :—

Causes of death by classes.