it, through the elimination of some of the zinc. No lead ore containing more than 20 p.c. of zinc was accepted, and for lead ores containing 4 p.c. of zinc or over the shipper was penalized according to the amount of the zinc present. A new schedule was announced in April, 1922, providing for payment for zinc in ores on a sliding scale running from 30 to 35 p.c. zinc.

Years.	Quantity ¹ .	Value.	Average price per pound.	Years.	Quantity ¹ .	Value.	Average price per pound.
	lb.	\$	ets.		lb.	\$	cts.
1911 1912 1913 1914 1915 1916	1,877,479 4,283,760 5,640,195 7,246,063 9,771,651 23,364,760 29,668,764		6.943 5.648 5.213 13.230 12.804	1918	35,083,175 32,194,707 39,863,912 53,089,356 56,290,000 60;416,240 98,909,077	2,362,448 3,057,961 2,471,310 3,217,536	7·338 7·671 4·655 5·716 6·607

25.—Production of Zinc in Canada, calendar years 1911-1924.

8.—Iron.1

The fact that iron ore is widely distributed in Canada has long been known, and extensive deposits have been discovered from time to time. The development of the iron-mining industry, however, has been retarded by the abundant supply of the higher grade ores of Wabana, Newfoundland, and of the Mesabi range of the state of Minnesota.

Nova Scotia.—The Wabana section of Newfoundland, containing the largest single deposit of iron ore in the world, is operated by the British Empire Steel Corporation. The probable reserves in that area have been estimated at 3,635,000,000 tons, and analysis has shown that the Wabana ore consists of an exceptionally high-grade hematite. Ore to the amount of 174,602 tons was shipped in 1924 to the blast-furnaces of the company at Sydney, where the proximity of the adjacent coal field favours the economical production of pig iron and steel. Development work carried on also at Torbrook, in Annapolis Co., indicates that the deposits there are very extensive. The ore is red hematite, containing a good percentage of iron rather high in phosphorus. An important iron ore field is the Arisaig district in Antigonish Co.

New Brunswick.—The most important deposits so far discovered are those in the Austin Brook district of Bathurst Co., where mining experts state that great masses of iron ore have been located.

Quebec.—It is estimated that there are many millions of tons of iron magnetite sands, containing a high percentage of iron, along the north shore of the St.Lawrence at Moisie, Mingan, Natashkwan and other places in Saguenay Co. The sands contain a high percentage of titanium, rendering the briquetted iron sands unfavourable for blast-furnace treatment. There are a number of deposits of bog iron ore in the St. Lawrence valley, remarkably free from sulphur and phosphorus. The bog iron ores were successfully used in charcoal blast-furnaces at Radnor Forges and Drummondville for many years. Iron ore deposits also exist along the Gatineau

¹ Estimated smelter recoveries, including for years 1916 to 1924 the actual zinc recovered at Trail, B.C.

A sketch of the iron and steel industry of Canada was given on pp. 452-456 of the 1922-1923 Year Book.