Death Rates by Age-Groups.—The death rate per 1,000 persons living in each group were shown in a table on p. 175 of the 1927-28 Year Book for the years 1921-24. The calculations were made on the assumption that the age constitution of the estimated population of the later years was the same as that of the ascertained population of the census year.

In this table there was indicated a declining death rate at the earlier ages, a stationary death rate between 35 and 55, and an increasing death rate after 55. While the Canadian period of observation is too short to establish these as general conclusions, the experience of other countries tends to confirm them as being common to the civilized countries of the world.

When the death-rate by sexes in various age-groups was considered it was evident that in most age-groups the female death-rate was lower than that of males, though there are significant exceptions. In 1924, the female death-rate was lower in the groups up to 20 and above 55. In the groups from 20-24 and 45-54 it was the same as that for males, but in the groups from 25 to 44 it was distinctly higher.

Adjusted Death Rate.—While the crude death rate gives the actual mortality per 1,000 of population, the differing age constitution of the population in different communities and the high mortality among infants and elderly people makes the crude death rate no true test of the relative expectation of life in such communities. Where the age constitution of a particular group is particularly favourable to low mortality, as, for example, in an army in peace time, the crude death rate will be lower than elsewhere.

When comparisons of the rates of mortality in several communities are made by age-groups as is done for the eight provinces in Table 21 on p. 162 of the Canada Year Book, 1925, the effects of differences in age constitution between these communities are eliminated, but by a rather cumbrous process, which does not bring together and express as a single figure the facts of the situation. It has therefore been considered desirable to adopt a particular community as a standard, and to find what the death rates of other communities would have been if the age and sex constitution of their population had corresponded to those of the community taken as a standard. The "standard" population chosen for this purpose in England and Wales and the United States is the "standard million", based on the age and sex distribution per million of the population was as follows:—

Age-groups.	Persons.	Males.	Females.
All ages.   Under 5 years.   5-9 years.   15-19 years.   15-19 years.   20-24 years.   25-34 years.   <	1,000,000 114,262 107,209 102,735 99,796 95,946 161,579 122,849 89,222 59,741 23,040	483,543 57,039 53,462 51,370 49,420 45,273 76,425 59,394 42,924 27,913 14,60	516, 457 57, 223 53, 747 51, 365 50, 376 50, 673 85, 154 463, 455 46, 298 31, 828
75 years and over	13,581	5,682	7.949