

27.—Infantile Mortality, by Provinces, together with the Rate per 1,000 Living Births, 1926-30 with Averages for 1921-25 and 1926-30.

A.—INFANT DEATHS.

Year.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Canada.*
Average 1921-25.....	151	1,139	1,165	1	5,916	1,394	1,789	1,327	621	1
Average 1926-30.....	122	934	1,039	10,518	5,091	1,031	1,559	1,195	571	22,060
1926.....	123	882	1,095	11,686	5,302	1,122	1,681	1,233	588	23,692
1927.....	113	1,028	1,006	10,739	4,812	1,021	1,575	1,110	606	22,010
1928.....	92	865	960	10,332	4,880	972	1,370	1,200	524	21,185
1929.....	150	960	1,090	9,810	5,203	1,005	1,571	1,310	575	21,674
1930.....	132	937	1,048	10,045	5,260	1,035	1,601	1,122	562	21,742

B.—INFANT DEATH RATE PER 1,000 LIVING BIRTHS.

Average 1921-25.....	76.8	93.7	105.0	1	82.7	83.7	83.0	85.8	60.6	1
Average 1926-30.....	70.6	84.8	100.7	127.1	74.1	71.8	73.3	75.5	55.2	93.3
1926.....	70.2	80.3	105.9	142.0	78.4	76.5	81.1	88.3	58.4	101.8
1927.....	66.6	92.3	96.0	129.3	71.1	72.2	74.9	74.5	60.1	94.0
1928.....	50.9	79.1	95.6	123.8	71.2	67.0	64.4	76.5	50.5	89.5
1929.....	89.3	89.8	106.5	120.5	76.0	70.6	73.3	77.4	55.4	92.1
1930.....	75.5	82.6	99.5	120.1	73.3	71.8	72.6	83.6	51.7	89.3

¹ Quebec not included in registration area prior to 1926.

* Exclusive of the Territories.

Infantile Mortality by Causes of Death.—Twenty principal causes of death accounted in the years 1926 to 1930 for between 92 and 93 p.c. of the infantile mortality experienced in the Dominion, as is shown in Table 28. It is noteworthy that four causes present at birth, *viz.*, premature birth, injuries at birth, congenital debility and congenital malformations, accounted for over 42 p.c. of the infant deaths of 1930. In the previous year 48.12 p.c. of all infants dying were less than one month old, and 33.63 p.c. less than one week old, as is shown in Table 29.

28.—Infantile Mortality in Canada by Principal Causes of Death, 1926-30.

Note.—Figures for the former registration area for the years 1921 to 1924 will be found at pp. 182-3 of the 1927-28 Year Book and for 1925 to 1927 at pp. 177-8 of the 1929 Year Book.

Cause of Death..	International List No.	Year.	Males.		Females.		Both Sexes.		Rate per 100,000 living births.
			No.	Rate per 100,000 living births.	No.	Rate per 100,000 living births.	No.	Per cent distribution by cause of death.	
Measles.....	7	1926	141	118	122	108	263	1.1	113
		1927	113	94	74	65	187	0.8	50
		1928	50	41	39	34	89	0.4	38
		1929	88	73	84	73	172	0.8	73
		1930	119	95	98	78	212	1.0	87
Scarlet fever.....	8	1926	13	11	12	11	25	0.1	11
		1927	23	19	12	11	35	0.2	15
		1928	12	10	13	11	25	0.1	11
		1929	10	8	10	9	20	0.1	8
		1930	17	14	18	11	30	0.1	12
Whooping cough.....	9	1926	358	299	415	368	773	3.3	332
		1927	336	278	304	268	640	2.9	273
		1928	230	189	239	207	469	2.2	193
		1929	231	191	223	195	454	2.1	183
		1930	345	276	341	237	686	3.2	232
Diphtheria	10	1926	24	20	23	20	47	0.2	20
		1927	24	20	24	21	48	0.2	20
		1928	22	18	26	23	48	0.2	20
		1929	22	18	21	18	43	0.2	18
		1930	29	23	20	17	49	0.2	20