


AVERAGE FLUCTUATIONS OF Q AND QW FOR 6 DISTRICTS SINCE BEFORE THE SUNSPOT MAXIMUM OF 1893 THROUGH 4 MAXIMA TO 1932 (SEE TABLE II FOR FIGURES)



9.—Precipitation in Crop Years and Temperature in the Growing Period at Certain Western Points, together with Quotients when Precipitation is Divided by Temperature, 1883-4-1932.

NOTE.—Quotients have in all cases been multiplied by 10.

EDMONTON.

Year.	P ₁₂	T ₃	Q	Year.	P ₁₂	T ₃	Q
1883-84.....	12.53	55.8	2.24	1909.....	15.15	57.1	2.66
85.....	16.06	55.8	2.44	10.....	10.82	57.1	1.90
86.....	8.30	57.2	1.45	11.....	20.90	56.8	3.68
87.....	13.00	53.9	2.42	12.....	20.18	57.9	3.48
88.....	21.84	54.4	4.02	13.....	20.25	56.4	3.60
89.....	7.37	57.6	1.28	14.....	21.47	57.4	3.74
90.....	16.97	56.5	3.01	15.....	21.95	54.9	4.00
91.....	22.40	56.9	3.94	16.....	17.19	55.0	3.12
92.....	13.76	54.8	2.51	17.....	19.34	56.4	3.43
93.....	19.94	56.2	3.55	18.....	16.52	55.0	3.00
94.....	14.33	58.7	2.44	19.....	15.66	57.0	2.75
95.....	15.50	56.7	2.73	20.....	21.01	56.2	3.74
96.....	16.06	57.4	2.80	21.....	16.62	57.4	2.90
97.....	15.15	58.6	2.58	22.....	10.88	57.8	1.90
98.....	11.97	59.0	2.03	23.....	18.86	57.5	3.28
99.....	14.44	55.1	2.62	24.....	16.43	57.4	2.87
1900.....	27.87	58.1	4.81	25.....	17.96	57.8	3.11
01.....	29.81	56.2	5.32	26.....	14.83	57.2	2.60
02.....	23.14	55.5	4.16	27.....	22.63	55.0	4.13
03.....	18.10	56.2	3.22	28.....	17.66	58.1	3.04
04.....	21.98	56.4	3.89	29.....	13.56	56.9	2.38
05.....	16.85	57.1	2.97	30.....	12.57	56.6	2.22
06.....	15.33	57.9	2.62	31.....	17.15	56.9	3.01
07.....	18.48	53.7	3.44	32.....	18.63	58.3	3.19
08.....	16.65	57.9	2.88				