16.—Weighted Quotients where Precipitation in Crop Year is Divided by Temperature in Growing Period, Expressed as a Percentage of Averages, 1886-1932—concluded.

NOTE.—For method of weighting see page 48.   Qw has in all cases been multiplied by 100.

Year.	QU'APPELLE.		MINNEDOSA.		WINNIPEG.	
	Q_w	Per cent of Average.	Q_w	Per cent of Average.	Q_w	Per cent of Average.
1886	217	67	293	100	301	91
1887	206	63	283	97	297	90
1888	256	79	283	97	284	86
1889	230	71	238	81	273	83
1890	291	89	265	90	306	93
1891	347	106	317	108	343	104
1892	334	102	284	97	370	112
1893	319	98	256	87	394	119
1894	240	74	246	85	339	103
1895	259	79	260	89	335	102
1896	297	91	302	103	370	112
1897	285	87	311	106	360	109
1898	262	80	281	96	331	100
1899	340	104	301	103	373	113
1900	298	91	266	91	315	95
1901	381	117	344	117	374	113
1902	436	134	364	124	354	107
1903	377	116	335	114	314	95
1904	394	121	358	122	344	104
1905	375	115	319	109	362	110
1906	391	120	325	111	374	113
1907	373	114	326	111	336	102
1908	344	106	311	106	325	98
1909	377	116	296	101	323	98
1910	374	115	251	85	329	100
1911	363	111	263	90	366	111
1912	362	111	318	109	365	110
1913	361	111	315	107	327	99
1914	393	121	290	99	319	97
1915	342	105	286	98	303	92
1916	366	112	318	109	355	108
1917	372	114	283	97	323	98
1918	344	106	250	85	298	90
1919	308	94	247	84	318	96
1920	295	90	264	90	335	102
1921	348	107	268	91	338	102
1922	379	116	321	110	332	101
1923	451	138	373	127	328	99
1924	361	111	323	110	274	83
1925	337	103	321	110	280	85
1926	286	88	244	83	266	81
1927	349	107	349	119	335	102
1928	369	113	357	122	379	115
1929	256	79	281	96	325	98
1930	240	74	244	83	321	97
1931	191	59	197	67	283	86
1932	236	72	249	85	300	91