accompanies the tables, the main object has been to attain the greatest degree of comparability possible over the whole period 1926-34. For this purpose it has been necessary to depart somewhat from the grouping of Tables 26 and 27. The cases in which this occurs are indicated on the chart by asterisks.

Total deaths in 1934 were the lowest they have been since statistics for all provinces of Canada were made possible by Quebec coming into the registration area in 1926. The influenza epidemic of 1929 accounted for an increase in the total, but the deaths from this cause have worked back to a normal level since that year. It is significant, however, that cancer and diseases of the arteries are increasingly important causes of death in spite of all efforts to control them. Suicides show an improvement from the high level reached in 1932. Violent deaths other than suicides, in which there was a very definite improvement between 1930 and 1933, showed an upward turn in 1934 as compared with the preceding year. A large proportion of this increase was contributed by automobile accidents.

26.—Deaths in Canada, by Principal Causes, 1929-34.

		<del></del>		<del></del>		<del>. :</del>	
Int. List No.1	Cause of Death.	1929.	1930.	1931.	1932.	1933.	1934.
7 8 9 10	Typhoid fever	467 619 440 755 980	451 521 397 964 737	421 167 253 748 646	339 330 197 555 398	291 170 157 552 239	293 188 226 875 232
16	Influenza <sup>2</sup>	7, 170 1 <b>5</b> 2	2,472 215	3,217 223	4,236 164	4,019 73	2,004 84
18 23 24-32 45-53 59 71	Lethargic or epidemic encephalitis Epidemic cerebro-spinal meningitis Tuberculosis, respiratory system <sup>2</sup> . Tuberculosis, other organs Cancer	118 341 6,443 1,329 8,792 1,208 693	88 294 6,581 1,494 9,273 1,146 740	77 225 6,204 1,412 9,578 1,244 716	76 139 5,870 1,296 10,024 1,343 728	58 109 5,664 1,275 10,653 1,287 736	47 84 5,290 1,141 10,581 1,321 612
82d 86 90–95	Cerebral hæmorrhage, embolism or thrombosis.  Paralysis without specified cause  Convulsions (under 5 years of age)  Diseases of the heart <sup>2</sup>	2,986 984 429 13,205	2,827 907 380 13,067	2,594 728 368 13,734	2,543 654 304 15,328	2,639 559 263 <sup>3</sup> 15,485	2,577 547 261 16,352
96, 97 99, 102 106 107–109 119–120	Diseases of the arteries?	5,940 471 8,441 4,910	6,560 443 7,338 6,013	5,957 469 7,011 5,158	6,798 437 7,045 3,735	6,950 367 6,487 3,395	7,379 380 6,530 3,730
121 122 130-132 137 140-150	Appendicitis.  Hernia, intestinal obstruction.  Nephritis.  Diseases of the prostate <sup>2</sup> .  Puerperal causes.	1,451 962 5,687 739 1,341	1,488 963 5,570 801 1,405	1,394 987 5,168 746 1,215	1,454 947 5,635 879 1,181	1,455 1,029 5,516 926 1,111	1,578 1,074 5,643 944 1,167
157 158–161 162 163–171	Congenital malformations Diseases of early infancy <sup>2</sup> Senility (old age) Suicides	1,466 9,144 2,505 835	1,475 8,974 2,334 1,010	1,427 9,019 2,225 1,004	1,349 7,932 2,192 1,024	1,374 7,337 2,037 922	1,361 6,936 1,882 927
173-198	Violent deaths (suicides excepted) Other specified causes <sup>2</sup> Totals, Specified Causes	$\frac{6,316}{15,020}$ $\frac{112,339}{112,339}$	6,468 14,919 108,315	6,168 12,914 103,417	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	5,294 12,546 100,975	5,542 12,857 100,645
199, <b>200</b>	Ill-defined diseases	1,176 113,515	991	1,100	1,007	993 101, <b>96</b> 8	937

¹ The numbers given in this column refer to the International List of Causes of Death, as revised in 1929 by the International Commission on the Classification of Diseases and Causes of Death. This classification, in its detailed, intermediate or abridged form, is accepted in almost all civilized countries.
² For these causes of death the comparability between the figures for the years 1929-30 and those for the years 1931-34 is not exact, owing to changes in classification.
² Revised since the publication of the 1934-35 Year Book.