23.—Infantile Mortality, by Provinces, together with the Rate per 1,000 Live Births, 1931-35, with Averages for 1921-25, 1926-30, and 1931-35.

A.—INFANT	

Year.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Canada.2
Averages, 1921-25	151	1,139	1,165	1	5,916	1,394	1,789	1,327	621	1
Averages, 1926-30	122	934	1,039	10,518	5,091	1,031	1,559	1,195	571	22,060
Averages, 1931-35	131 128	840 914	857 944	7,756 9,443	3,962 4,833	835 924	1,261	998	464 514	17,10 20,36
1932	132	849	774	7.744	4,133	836	1.321	997	477	17,263
1933	132 118	791	821	7,270	3,804	844	1,231	966	439	16,28
1934	130	807	878	7,388	3,523	734	1,093	891	426	15,87
1935	145	838	866	6,939	3,515	837	1,194	936	460	15,73

B .- INFANT DEATH RATES PER 1,000 LIVE BIRTHS.

- 24	- 1	1	1			1		1	
77	94	105	1	83	84	83	86	61	1
71	85	101	127	74	72	73	75	55	93
67	73	82	98	61	61	62	60	46	75
68	79	87	113	70	64	69	69	49	85
65	73	72	94	62	59	63	59	47	73
61	71	82	95	60	63	61	60	46	73
67	71	86	97	57	55	55	55	43	72
72	72	83	92	56	63	61	58	46	71
	65 61 67	71 85 67 73 68 79 65 73 61 71 67 71	71 85 101 67 73 82 68 79 87 65 73 72 61 71 82 67 71 86	71 85 101 127 67 73 82 98 68 79 87 113 65 73 72 94 61 71 82 95 67 71 86 97	71 85 101 127 74 67 73 82 98 61 68 79 87 113 70 65 73 72 94 62 61 71 82 95 60 67 71 86 97 57	71 85 101 127 74 72 67 73 82 98 61 61 68 79 87 113 70 64 65 73 72 94 62 59 61 71 82 95 60 63 67 71 86 97 57 55	71 85 101 127 74 72 73 67 73 82 98 61 61 62 68 79 87 113 70 64 69 65 73 72 94 62 59 63 61 71 82 95 60 63 61 67 71 86 97 57 55 55	71 85 101 127 74 72 73 75 67 73 82 98 61 61 62 60 68 79 87 113 70 64 69 69 65 73 72 94 62 59 63 59 61 71 82 95 60 63 61 60 67 71 86 97 57 55 55 55	71 85 101 127 74 72 73 75 55 67 73 82 98 61 61 62 60 46 68 79 87 113 70 64 69 69 49 65 73 72 94 62 59 63 59 47 61 71 82 95 60 63 61 60 46 67 71 86 97 57 55 55 55 43

¹ Quebec was not included in the registration area prior to 1926.

Infantile Mortality by Causes of Death.—Twenty-one principal causes of death accounted in the years 1926 to 1935 for between 91 and 92 p.c. of the infantile mortality experienced in the Dominion, as is shown in Table 24. It is noteworthy that four causes present at birth, viz., premature birth, injury at birth, congenital debility and congenital malformations, accounted for nearly 44 p.c. of the infant deaths of 1935. This percentage shows a definite increase in the years covered. In 1926 it was 41.4 and in 1930 42.3, and since the total number of infant deaths has decreased by no less than 34 p.c. in the interval since 1926, great improvement in the post-natal care of infants is indicated. In the years 1934 and 1935, 49.0 p.c. and 49.2 p.c., respectively, of all infants who died were less than one month old, and 34.9 p.c. and 35.3 p.c., respectively, were less than one week old, as is shown in Table 25.

24.—Infantile Mortality in Canada by Principal Causes of Death, 1926, 1934-35.

Note.—Figures for the former registration area for the single years 1921 to 1924 will be found at pp. 182-183 of the 1927-28 Year Book, and for 1925 to 1927 at pp. 177-178 of the 1929 Year Book. Figures for the whole of Canada for the years 1927 and 1928 will be found at pp. 138-140 of the 1932 Year Book, for 1929 and 1930 at pp. 177-178 of the 1933 Year Book, and for 1931, 1932 and 1933 at pp. 202-203 of the 1934-35 Year Book.

Inter- national List No.	Cause of Death.	Year.	1	Numbers.		Rat L	Percent- age Distribu-		
			Males.	Fe- males.	Both.	Males.	Fe- males.	Both.	tion by Cause of Death.
7	Measles	1926 1934	141 43	122 30	263 73	118 38	108 28 71	113 33	1·1 0·5
8	Scarlet fever	1935 1926 1934 1935	104 13 6 8	77 12 1	181 25 7 15	92 11 5	11 11 1	82 11 3	1 · 2 0 · 1 1 0 · 1
9	Whooping-cough	1926 1934 1935	358 296 318	415 309 281	773 605 599	299 261 281	368 286 260	332 273 270	3·3 3·8 3·8

For footnotes, see end of table, p. 182.

² Exclusive of the Territories.