20.—Death Rates per 100,000 Population in Canada, by Principal Causes, 1933-37.

Int. List No.	Cause of Death.	1933.	1934.	´1935.	1936.	1937.
1,2	Typhoid fever	2.7	2.7	2.5	2-3	3.0
7	Measles	1.6	1.7	4.5	3.4	7.5
	Scarlet fever	1.5	2.1	2-2	2.2	2.4
.9	Whooping-cough	5-2	8.1	8.2	5.4	6.9
	Diphtheria	2.2	2.1	2.4	2.3	3.3
	Influenza	37.7	18.5	31.1	28.3	47-4
16	Poliomyelitis and polioencephalitis (acute)	0.7	0.8	0.6	0.9	1.8
17	Lethargic or epidemic encephalitis	0·5 1·0	0·4 0·8	0·5 1·0	0·5 0·9	0·5 0·8
18 23	Epidemic cerebrospinal meningitis	53·1	48.9	50·1	50.2	0.8 49.5
	Tuberculosis, respiratory system	12-0	10.6	10.4	11.2	10.6
45-63	Cancer	99.9	97.9	102.2	106.2	107.7
59	Disbetes mellitus	12.1	12.2	13.4	13-1	14.0
71	Ausmia	6.9	5.7	6.0	5-9	5.6
	Cerebral hæmorrhage, embolism or thrombosis,		23.8	19.3	17-2	15.2
	Paralysis without specified cause	5.2	5.1	3.8	3.3	2.9
86	Convulsions (under 5 years of age)	2.5	2.4	2.1	1.8	1.8
90-95	Diseases of the heart	145.2	151-3	147-1	149-1	151.6
96, 97, 99, 102	Diseases of the arteries	65.2	68.3	76.0	82.7	86.5
	Bronchitis	3.4	3.5	3.3	3-1	3.0
107-109	Pneumonia	60-8	60-4	67-9	66-4	69-6
119, 120	Diarrhœa and enteritis	31.8	34.5	25.8	21.6	38-0
121	Appendicitis	13.6	14-6	13.7	13.0	12.7
122		9.6	9.9	10-3	9.5	9.7
130-132	Nephritis	51.7	52.2	56-6	58∙1	58-8
137	Diseases of the prostate	8.7	8.7	10.0	10-5	11.8
140-150	Puerperal causes	10-≰	10-8	10-0	11-2	9-6
157	Congenital malformations	12.9	12-6	13.0	13 - 1	13.3
158-161		68.8	64-2	63 0	60.0	59-8
162		19-1	17.4	17.7	15.4	15.7
163-171	Saicides	8.6	8.6	8.3	8.4	8.8
173-198	Violent deaths (suicides excepted)	49 6	51.3	54.9	59.3	57-4
	Other specified causes	117-6	118-9	122 · 6	129 - 1	131-4
	Totals, Specified Causes	946-6	931.0	959-7	965-5	1,017-9
199, 200	Ill-defined diseases	9.3	8-7	7.0	6.5	7.0
	Totals, Death Rates per 100,000 Population	955 - 9	939-7	966-6	571-9	1,024-1

For footnote, see end of Table 19, p. 136.

Deaths in Canadian Cities.—Table 21 gives the numbers of deaths in Canadian cities and towns of 10,000 population or over for the years 1933-37, together with averages, 1926-30 and 1931-35. Deaths in Canada as a whole declined steadily for the period 1931-34, but for 1935, 1936, and 1937 there were substantial increases. The figure for the latter year was 113,824, over 9,000 more than for 1931. The total deaths of the 67 cities listed in Table 21 show a slightly increased proportion to population for the five-year period 1931-35 as compared with 1926-30. For 1932, which marked the depth of the economic depression, the deaths in these cities increased, thus going against the general trend for Canada; for other years, however, the general trend was followed.

Deaths by Place of Residence.—The Vital Statistics Branch of the Dominion Bureau of Statistics, has, at the time of going to press, a report in process of compilation showing deaths by places of residence, (see p. 116).