

deaths of 1937. In 1926 it was 41·4 and in 1930, 42·3, and since the decline in rate of infant deaths has decreased by nearly 30 p.c. in the interval since 1926, great improvement in the post-natal care of infants is indicated. In the years 1936 and 1937, 50·7 p.c. and 45·1 p.c., respectively, of all infants who died were less than one month old, and 37·2 p.c. and 32·5 p.c., respectively, were less than one week old, as is shown in Table 25.

24.—Infantile Mortality in Canada by Principal Causes of Death, 1926, 1936-37.

Note.—Figures for the former registration area for the single years 1921 to 1924 will be found at pp. 182-183 of the 1927-28 Year Book, and for 1925 to 1927 at pp. 177-178 of the 1929 Year Book. Figures for the whole of Canada for the years 1927 and 1928 will be found at pp. 138-140 of the 1932 Year Book, for 1929 and 1930 at pp. 177-178 of the 1933 Year Book, for 1931, 1932, and 1933 at pp. 202-203 of the 1934-35 Year Book, for 1934 at pp. 176-177 of the 1936 Year Book, and for 1935 at pp. 182-183 of the 1938 Year Book.

| International List No. | Cause of Death. | Year | Numbers. | | | Rates per 100,000 Live Births. | | | Percentage Distribution by Cause of Death. |
|------------------------|---|------|----------|----------|-------|--------------------------------|----------|-------|--|
| | | | Males. | Females. | Both. | Males. | Females. | Both. | |
| 7 | Measles..... | 1926 | 141 | 122 | 263 | 118 | 108 | 113 | 1·1 |
| | | 1936 | 66 | 49 | 115 | 53 | 46 | 52 | 0·8 |
| | | 1937 | 158 | 110 | 268 | 140 | 103 | 122 | 1·6 |
| 8 | Scarlet fever..... | 1926 | 13 | 12 | 25 | 11 | 11 | 11 | 0·1 |
| | | 1936 | 7 | 4 | 11 | 6 | 4 | 5 | 0·1 |
| | | 1937 | 6 | 5 | 11 | 5 | 5 | 5 | 0·1 |
| 9 | Whooping-cough..... | 1926 | 358 | 415 | 773 | 299 | 368 | 322 | 3·3 |
| | | 1936 | 203 | 189 | 392 | 179 | 177 | 178 | 2·7 |
| | | 1937 | 216 | 269 | 485 | 191 | 251 | 220 | 2·9 |
| 10 | Diphtheria..... | 1926 | 24 | 23 | 47 | 20 | 20 | 20 | 0·2 |
| | | 1936 | 8 | 3 | 11 | 7 | 3 | 5 | 0·1 |
| | | 1937 | 10 | 14 | 24 | 9 | 13 | 11 | 0·1 |
| 11 | Influenza ¹ | 1926 | 576 | 274 | 950 | 481 | 331 | 408 | 4·0 |
| | | 1936 | 344 | 232 | 576 | 304 | 217 | 261 | 4·0 |
| | | 1937 | 545 | 394 | 939 | 482 | 368 | 426 | 5·6 |
| 15 | Erysipelas..... | 1926 | 51 | 50 | 101 | 43 | 44 | 43 | 0·4 |
| | | 1936 | 42 | 27 | 69 | 37 | 25 | 31 | 0·5 |
| | | 1937 | 13 | 9 | 22 | 11 | 8 | 10 | 0·1 |
| 16 | Poliomyelitis and polio-encephalitis (acute)..... | 1926 | 6 | 3 | 9 | 5 | 3 | 4 | 0·3 |
| | | 1936 | 1 | 3 | 4 | 1 | 3 | 2 | 1 |
| | | 1937 | 7 | 2 | 9 | 6 | 2 | 4 | 0·1 |
| 18 | Epidemic cerebro-spinal meningitis..... | 1926 | 33 | 24 | 57 | 28 | 21 | 24 | 0·2 |
| | | 1936 | 11 | 4 | 15 | 10 | 4 | 7 | 0·1 |
| | | 1937 | 12 | 12 | 24 | 11 | 11 | 11 | 0·1 |
| 23-32 | Tuberculosis ¹ | 1926 | 131 | 102 | 233 | 109 | 90 | 100 | 1·0 |
| | | 1936 | 93 | 65 | 158 | 82 | 61 | 72 | 1·1 |
| | | 1937 | 85 | 64 | 149 | 75 | 60 | 68 | 0·9 |
| 34 | Syphilis..... | 1926 | 68 | 60 | 138 | 57 | 53 | 55 | 0·5 |
| | | 1936 | 118 | 59 | 207 | 104 | 83 | 94 | 1·4 |
| | | 1937 | 121 | 81 | 202 | 107 | 76 | 92 | 1·2 |
| 86 | Convulsions..... | 1926 | 263 | 177 | 440 | 218 | 157 | 189 | 1·9 |
| | | 1936 | 107 | 55 | 162 | 94 | 51 | 74 | 1·1 |
| | | 1937 | 83 | 56 | 139 | 73 | 52 | 63 | 0·8 |
| 106 | Bronchitis..... | 1926 | 90 | 60 | 150 | 75 | 53 | 64 | 0·6 |
| | | 1936 | 39 | 37 | 76 | 34 | 35 | 34 | 0·5 |
| | | 1937 | 34 | 32 | 66 | 30 | 30 | 30 | 0·4 |
| 107-109 | Pneumonia..... | 1926 | 1,410 | 1,077 | 2,487 | 1,176 | 954 | 1,069 | 10·5 |
| | | 1936 | 967 | 783 | 1,750 | 854 | 731 | 794 | 12·0 |
| | | 1937 | 1,105 | 809 | 1,914 | 977 | 755 | 869 | 11·5 |
| 116-118 | Diseases of the stomach..... | 1926 | 156 | 126 | 282 | 130 | 112 | 121 | 1·2 |
| | | 1936 | 70 | 40 | 110 | 62 | 37 | 50 | 0·8 |
| | | 1937 | 52 | 43 | 95 | 46 | 40 | 43 | 0·6 |

¹ For these causes the comparability between the figures for the year 1926 and those for the years 1936-37 is not exact, owing to changes in classification.

² Less than one-tenth of one per cent.