

**6.—Diagnoses of Patients on the Books of Reporting Tuberculosis Hospitals, by Class, and by Age and Sex, 1958**

NOTE.—Figures include patients in 82 tuberculosis hospitals and also patients in tuberculosis units of general hospitals.

Diagnosis	0-9		10-19		20-29		30-39	
	M.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Respiratory tuberculosis.....	447	429	387	506	737	730	782	680
With occupational disease of lung.....	—	—	—	—	1	1	—	—
Minimal pulmonary.....	65	55	112	163	149	189	145	169
Moderately advanced pulmonary.....	35	28	106	147	348	314	325	273
Far advanced pulmonary.....	12	8	58	76	195	189	281	210
Other and unspecified pulmonary.....	22	27	5	14	5	2	7	8
Pleurisy with or without effusion.....	12	12	42	39	36	26	24	17
Primary infection.....	287	285	61	62	3	5	—	1
Other.....	16	14	3	5	—	4	—	2
Tuberculosis, other forms.....	58	57	59	53	47	59	58	47
	40-49		50-59		60+		Total <sup>1</sup>	
	M.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Respiratory tuberculosis.....	834	459	972	317	1,372	470	5,655	3,660
With occupational disease of lung.....	—	1	3	—	9	—	14	3
Minimal pulmonary.....	104	97	116	52	131	70	844	815
Moderately advanced pulmonary.....	335	192	379	127	598	196	2,172	1,299
Far advanced pulmonary.....	369	159	460	128	610	191	2,028	976
Other and unspecified pulmonary.....	5	2	4	1	7	2	55	58
Pleurisy with or without effusion.....	21	7	9	6	15	9	161	116
Primary infection.....	—	—	—	—	—	—	358	361
Other.....	—	1	1	3	2	2	23	32
Tuberculosis, other forms.....	40	31	29	21	32	20	331	294

<sup>1</sup> Includes age groups not known.

**Facilities.**—Table 7 shows the prevalence of special facilities in public and federal hospitals distributed by province. Nationally, over 75 p.c. of these hospitals were equipped with radiology, clinical laboratory and out-patient facilities, but fewer than 50 p.c. with physiotherapy facilities. Among other factors, the type of service offered in a hospital determines to some extent the facilities provided. For example, about 84 p.c. of the general hospitals reported radiology, clinical laboratory and out-patient facilities, and nearly half of these hospitals provided physiotherapy facilities. On the other hand, only just over one of every three mental institutions had clinical laboratory, physiotherapy and out-patient facilities, although about two of every three had radiology facilities. Among tuberculosis institutions, slightly more than three of every four were equipped with radiology facilities, about 85 p.c. with clinical laboratory, and only about one of every three with physiotherapy and out-patient facilities. Fewer than one-third of "other" hospitals provided radiology, clinical laboratory and out-patient facilities, while slightly more than one-third were equipped for physiotherapy.