Diagnoses of Patients in Mental Institutions and Tuberculosis Sanatoria.— Tables 6 and 7 summarize the most recent data available on diagnoses according to age and sex of patients on the books of mental institutions and tuberculosis sanatoria, including patients in related units located in general hospitals.

Of the 71,252 mentally ill patients on record at Dec. 31, 1958, 64.5 p.c. had been diagnosed as psychotic, 31.6 p.c. as mentally defective or otherwise affected by character, behaviour and intelligence disorders, and 2.5 p.c. as psychoneurotic. More men than women had been institutionalized for disorders of character, behaviour and intelligence, with the largest frequencies of both sexes occurring in the 10-19 age group. Similarly, more men than women (except of persons 60 years of age or over) had been diagnosed as psychotic. However, in all age groups, nearly twice as many women as men had required hospitalization for conditions diagnosed as psychoneurotic.

6.—Diagnoses of Patients on the Books of Reporting Mental Hospitals, by Age and Sex, 1958

Note.—Figures include patients in 78 mental hospitals and also patients in psychiatric and mental units of general hospitals.

Diagnosis	0-9		10-19		20-29		30-39	
	M,	F.	M.	F.	M.	F.	M.	F.
Psychoses Schizophrenia. Manic-depressive reaction. Of old age. Other.	No. 17 13 — 4	No. 8 6 - 2	No. 262 198 15 -49	No. 213 140 15 - 58	No. 2,043 1,664 90 - 289	No. 1,353 1,001 135 — 217	No. 4,022 3,156 225 641	No. 3,132 2,339 328 465
Psychoneuroses Neurotio-depressive reaction Anxiety reaction Other	1111	Ξ	13 7 2 4	36 11 5 20	78 25 24 29	149 68 21 60	107 42 26 39	267 136 39 92
Disorders of Character, Behaviour and Intelligence. Alcoholism. Mental deficiency. Other.	1,172 1,074 98	961 	3,473 1 3,214 258	2,509 	2,490 18 2,160 312	1,974 11 1,712 251	1,987 94 1,574 319	1,607 21 1,313 273
Non-psychiatric and Unreported Diag- noses	19	21	41	34	56	26	71	48
	40-49		50-59		60+		Total ¹	
	М.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Psychoses Schizophrenia. Manic-depressive reaction. Of old age. Other.	4,886 3,406 387 8 1,085	4,180 2,652 567 9 952	5,186 2,939 490 68 1,689	4,330 2,325 597 45 1,363	7,995 3,058 739 2,037 2,161	8,246 2,878 1,081 2,434 1,853	24,458 14,463 1,947 2,117 5,931	21,506 11,365 2,733 2,492 4,916
Psychoneuroses. Neurotic-depressive reaction. Anxiety reaction. Other.	122 48 18 56	243 110 36 97	126 50 17 59	217 81 18 118	142 78 13 51	273 126 25 122	589 250 101 238	1,185 536 144 509
Disorders of Character, Behaviour and Intelligence. Alcoholism. Mental deficiency. Other.	1,494 137 1,133 224	1,350 34 1,108 208	999 102 738 159	875 16 724 135	892 85 665 142	737 17 613 107	12,526 439 10,572 1,515	10,025 99 8,621 1,305
Non-psychiatric and Unreported Diag-	90	51	98	64	175	162	552	407

¹ Includes age groups not known.