

13.—Diagnoses of Patients on the Books of Reporting Mental Hospitals, by Age and Sex, 1959 —concluded

Diagnosis	0-9		10-19		20-29		30-39	
	M.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Psychoneuroses	—	—	16	35	79	174	140	289
Neurotic-depressive reaction.....	—	—	5	19	32	86	62	142
Anxiety reaction.....	—	—	—	3	23	22	31	35
Other.....	—	—	11	13	24	66	47	112
Disorders of Character, Behaviour and Intelligence	1,333	1,635	3,771	2,620	2,581	2,626	2,110	1,681
Alcoholism.....	—	—	1	—	23	4	124	26
Mental deficiency.....	1,223	953	3,453	2,330	2,243	1,756	1,627	1,352
Other.....	110	82	317	290	315	266	359	303
Non-psychiatric and Unreported Diagnoses	19	22	45	32	55	17	72	45
	40-49		50-59		60+		Total ¹	
	M.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Psychoses	4,843	4,148	5,124	4,458	8,140	8,394	24,502	21,713
Schizophrenia.....	3,396	2,684	2,945	2,417	3,112	2,922	14,534	11,529
Manic-depressive reaction.....	396	516	494	609	778	1,057	2,021	2,688
Of old age.....	6	8	73	42	2,102	2,496	2,185	2,548
Other.....	1,045	940	1,612	1,390	2,148	1,919	5,762	4,948
Psychoneuroses	141	265	143	235	160	288	680	1,288
Neurotic-depressive reaction.....	57	126	59	91	84	136	299	602
Anxiety reaction.....	22	29	19	15	15	26	111	130
Other.....	62	110	65	129	61	126	270	556
Disorders of Character, Behaviour and Intelligence	1,556	1,368	1,061	931	926	770	13,351	10,442
Alcoholism.....	148	32	127	29	98	17	521	108
Mental deficiency.....	1,167	1,117	766	748	680	633	11,171	8,900
Other.....	241	219	168	154	148	120	1,659	1,434
Non-psychiatric and Unreported Diagnoses	86	41	106	63	257	120	641	430

¹ Includes age groups not known.

Among patients reported as on the books of tuberculosis institutions at the end of 1960, the peak frequency for men (1,199) occurred in the 60 years or over age group, and for women (480) in the 20-29 age group. More men than women were listed in each age group except those encompassing ages 10-29. Compared with 1,199 men, only 387 women aged 60 or over were carried on the books of tuberculosis sanatoria and units at Dec. 31, 1960; in contrast, the peak frequency for women aged 20-29 (480) was closely approximated in number by male patients (497).

14.—Diagnoses of Patients on the Books of Reporting Tuberculosis Hospitals, by Age and Sex, 1960

NOTE.—Figures include patients in 70 tuberculosis hospitals and patients in tuberculosis units in general hospitals but exclude tubercular patients in mental hospitals.

Diagnosis	0-9		10-19		20-29		30-39	
	M.	F.	M.	F.	M.	F.	M.	F.
	No.	No.	No.	No.	No.	No.	No.	No.
Respiratory Tuberculosis	353	339	398	311	448	435	519	427
With occupational disease of lung.....	—	—	—	—	—	—	1	—
Minimal pulmonary.....	52	56	70	104	97	111	75	94
Moderately advanced pulmonary.....	21	19	77	89	189	179	240	177
Far advanced pulmonary.....	4	4	46	36	134	120	193	144
Other and unspecified pulmonary.....	17	20	7	9	2	4	3	3
Pleurisy with or without effusion.....	7	3	30	26	26	17	7	7
Primary infection.....	246	220	69	44	—	4	—	—
Other.....	6	17	9	3	—	—	—	2
Tuberculosis, Other Forms	37	41	29	36	49	45	62	36