

**12.—Revenues and Expenditures of Reporting Public Hospitals per Patient-Day
(excluding Newborn), by Province, 1962—concluded**

Province or Territory and Type of Hospital	Hospitals Reporting	Total Revenue	Expenditures				
			Gross Salaries and Wages	Medical and Surgical Supplies	Drugs	Other	Total
	No.	\$	\$	\$	\$	\$	\$
British Columbia—							
General.....	86	24.21	17.36	0.80	0.98	5.89	25.03
Allied Special—							
Chronic, convalescent and rehabilitation	3	16.80	12.48	0.14	0.17	4.58	17.37
Maternity.....	1	28.87	22.19	0.82	0.96	6.45	30.42
Other.....	9	14.28	13.92	0.12	0.29	7.82	22.15
Mental.....	8	7.28	4.44	—	0.32 ¹	2.52	7.28
Tuberculosis.....	2	16.82	12.31	0.72	0.28	3.51	16.82
Yukon Territory—							
General.....	2	30.38	18.18	0.66	1.91	11.77	32.52
Northwest Territories—							
General.....	3	18.96	11.64	0.32	0.58	9.65	22.19

¹ Medical and surgical supplies are included with drugs.

Diagnoses of Patients in Institutions for Tuberculosis and for Psychiatric Disorders.—There were 4,824 patients with respiratory tuberculosis on the books of tuberculosis institutions on Dec. 31, 1962, and 63 p.c. of them were males. Except for the age groups under 20 years, males outnumbered females in all age groups. The peak frequency for men (963) occurred in the 60+ age group, and for women (554) in the 20-39 age group.

Of the 75,300 psychiatric patients reported at Dec. 31, 1962, 34 p.c. suffered from schizophrenia, the most frequent form of psychosis (insanity); 25 p.c. suffered from other forms of psychosis; and 31 p.c. were mentally deficient. In most diagnostic classes males outnumbered females; exceptions were certain psychoses, neuroses and drug addiction.

13.—Diagnoses of Patients on the Books of Reporting Tuberculosis Hospitals, by Age Group and Sex, as at Dec. 31, 1962

NOTE.—Figures include patients in 65 tuberculosis hospitals and patients in tuberculosis units in general hospitals but exclude tubercular patients in mental hospitals.

Diagnosis	Age Group					Total ¹
	0-14	15-19	20-39	40-59	60 +	
	NUMBERS					
Respiratory tuberculosis.....	M. 334	113	679	953	963	3,047
With occupational disease of lung.....	F. 333	121	554	428	339	1,777
Other.....	M. —	—	—	23	42	65
Minimal pulmonary.....	F. —	—	—	—	—	—
Moderately advanced pulmonary.....	M. 36	29	117	123	106	411
Far advanced pulmonary.....	F. 47	39	136	86	62	370
Other.....	M. 16	45	283	344	384	1,074
Pleurisy with or without effusion.....	F. 22	48	248	185	141	645
Primary infection.....	M. 6	12	241	448	421	1,131
Other.....	F. 3	17	142	140	133	436
Tuberculosis, other forms.....	M. 6	3	8	2	4	23
Other.....	F. 12	3	7	3	—	25
Pleurisy with or without effusion.....	M. 18	15	28	12	5	78
Primary infection.....	F. 7	7	18	12	3	47
Other.....	M. 237	8	2	—	—	247
Tuberculosis, other forms.....	F. 232	7	3	—	—	242
Other.....	M. 15	1	—	1	1	18
Tuberculosis, other forms.....	F. 10	—	—	2	—	12
Other.....	M. 34	11	54	54	33	186
Tuberculosis, other forms.....	F. 40	19	63	34	23	179

¹ Includes ages not known.