from year to year. As is evident from Table 2, pp. 245-247, many cities and towns have maintained consistently low rates as compared with the national rate or the rate for the province in which they are situated.

Causes of Infant Deaths.—In 1963 almost 70 p.c. of the infant deaths were caused by immaturity, congenital malformations, pneumonia, postnatal asphyxia and atelectasis, and injury at birth. Immaturity was the underlying cause of 2,348 and was an added complication in 2,972 others. Congenital malformations accounted for 2,068 fatalities, pneumonia for 1,506, postnatal asphyxia for 1,368 and injury at birth for 1,232. Rates for congenital malformations and birth injuries decreased in 1963.

21.-Infant Mortality and Rates per 100,000 Live Births, by Cause, 1961-63

Inter- national List No.	Cause of Death	Numbers of Deaths			Rates per 100,000 Live Births		
		1961	1962	1963	1961	1962	1963
001-019 020-029 045-048 050 052 055 056 057 085 140-239 273 340 391, 392 470-475 480-483 500-502 571 762 750-769 760, 761 763 764 765-768 769 770 771 772 774-776 E810-E825 E900-E904 E921, E922	Tuberculosis Syphilis Dysentery Scarlet fever Erysipelas Diphtheria Whooping cough Meningococal infections Measles Neoplasms Diseases of thymus gland Mental deficiency Meningitis (non-meningococcal) Otitis media Acute upper respiratory infections Influensa Pneumonia (4 weeks and over) Bronchitis Gastritis and duodenitis Hernia and intestinal obstruction Gastro-enteritis and colitis Chronic enteritis, and ulcerative colitis Congenital malformations Injury at birth Postnatal asphyxia and atelectasis Pneumonia of newborn (under 4 weeks) Diarrheza of newborn (under 4 weeks) Other infections of newborn Antenatal toxacmia Erythroblastosis Hæmorrhagic disease of newborn Nutritional maladjustment Ill-defined and unknown causes Motor vehicle accidents Accidents caused by fire Inhalation and ingestion of food or other object Accidental mechanical suffocation Other accidental and violent deaths Other specified causes	8 1 1 14 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 3 3 7 1 1 19 19 34 4 37 7 19 64 66 45 99 4 66 45 97 1,232 84 2,230 4 1,338 1,475 408 46 66 31 31 31 31 31 477 51 473	5 3 3 1 — — — — — — — — — — — — — — — — —	2 3 3 5 6 8 6 8 6 8 6 8 6 8 6 8 6 8 6 8 6 8 6	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	Totals, All Causes	12,940	12,941	12,270	2,720	2,755	2,634

Age at Death.—Of the 12,270 infants who died within a year of their birth, 8,411, or over 68 p.c., were less than one month old—4,904 during the first day of life, 2,569 from the second to the seventh day, and 938 during the three following weeks.