Perinatal Mortality.—'Perinatal' mortality—the combined total of stillbirths and deaths of live-born infants occurring 'around' the natal period—is a relatively new vital statistics concept. Since such deaths frequently have the same underlying causes, associated with pregnancy or delivery, regardless of whether they occur before or after delivery, perinatal deaths are generally considered as including the combined total of stillbirths occurring after at least 28 weeks pregnancy and deaths of live-born infants who fail to survive the first week of life.

In 1964 there were 12,539 such 'deaths', of which 5,520 were stillborn and 7,019 liveborn but failed to survive one week, with a national rate of 27.4 such deaths for every 1,000 total deliveries. This perinatal rate has declined slowly but steadily from 65.2 in 1921 to 27.4 in 1964.

Subsection 3.—Maternal Mortality

As indicated in Table 1, pp. 236-237, the number of mothers who die in pregnancy and childbirth has been greatly reduced during the past two decades, reaching an all-time low of 137 in 1964. Since 1951 the rate of maternal mortality per 10,000 births has been under 10 and since 1959 it has been under five. Despite this improvement, Canada's maternal death rate (3.0 in 1964) is higher than those for several other countries (see p. 276). Mortality among unmarried mothers is higher than among married mothers.

Causes of Maternal Deaths.—Table 24 shows the main causes of maternal deaths during the years 1962-64.

Inter- national List No.	Cause of Death	Numbers of Deaths			Rates per 100,000 Live Births		
		1962	1963	1964	1962	1963	1964
640,641	Complications of Pregnancy	52	43	35	11	9	8
010,011	pregnancy	3	6	3	1	1	1
642	Toxæmias of pregnancy	23	21	17	5	5	4
645	Ectopic pregnancy	6	8	6	ĩ	5 2	1
646-649	Other complications of pregnancy	20	8	9	4	2	2
	Abortion	24	27	22	5	6	5 2 3
650,652	Abortion without mention of sepsis	9	6	10	2 3	1	2
651	Abortion with sepsis	15	21	12	3	5	3
670	Complications of Delivery	78	67	48	17	14	11
670	Delivery complicated by placenta prævia or antepartum hæmorrhage	21	20	6	4	4	1
671	Delivery complicated by retained placenta	3	5	4	1	1	ĺ
672	Delivery complicated by other postpartum	9099W	50 1	8595	2500	900	
673,674	hæmorrhage Delivery complicated by abnormality of bony	21	6	15	4	1	8
010,011	pelvis or malposition of fœtus	6	8	5	1	2	1
675	Delivery complicated by prolonged labour of				**	2.70	
	other origin	5	4	1	1	1	-0
676,677 678	Delivery with laceration or other trauma Delivery with other complications of child-	8	10	11	2	2	2
	birth	14	14	6	3	3	1
	Complications of the Puerperium	37	28	32	8	6	2
680	Puerperal urinary infection without other sepsis	S-70	100	1	-	-	- i
681	Sepsis of childbirth and the puerperium	13	4	5	3	1	1
682-684	Puerperal phlebitis, thrombosis, pyrexia, pul-	10					
	monary embolism	12	9	13	3	2	3
685,686	Puerperal eclampsia and toxemia	.1	5	1		1	
687-689	Other	11	10	12	2	2	8
	Totals, All Puerperal Causes	191	165	137	41	35	30