

well represented in the north-central portions. Engelmann spruce and alpine fir from the Subalpine Region together with white birch are important constituents in the northern parts. The white spruce, though primarily boreal in affinity, is also present here. Extensive prairie communities of bunch-grasses and forbs are found in many of the river valleys.

*Coast Forest Region.*—This is part of the Pacific Coast forest of North America. Essentially coniferous, it consists principally of western red cedar and western hemlock, with abundant Sitka spruce in the north and with the addition of Douglas fir in the south. Amabilis fir and yellow cedar occur widely and, together with mountain hemlock and alpine fir, are common toward the timber-line. Western white pine is found in the southern parts and western yew is scattered throughout. Broadleaved trees, such as black cottonwood, red alder and broadleaf maple, have a limited distribution. Arbutus and Garry oak, species whose centres of population lie southward in the United States, occur in Canada only on the southeast coast of Vancouver Island and the adjacent islands and mainland.

*Acadian Forest Region.*—Over the greater part of the Maritime Provinces, exclusive of Newfoundland, there is a forest closely related to the Great Lakes-St. Lawrence Region and, to a lesser extent, to the Boreal Region. Red spruce is a characteristic though not exclusive species and associated with it are balsam fir, yellow birch and sugar maple, with some red pine, white pine and hemlock. Beech was once an important forest constituent but the beech bark disease has drastically reduced its abundance in Nova Scotia, Prince Edward Island and southern New Brunswick. Other species of wide distribution are the black and the white spruces, red oak, white elm, black ash, red maple, white birch, wire birch and the poplars. Eastern white cedar, though present in New Brunswick, is extremely rare elsewhere and jack pine is apparently absent from the upper St. John Valley and the western half of Nova Scotia.

*Columbia Forest Region.*—A large part of the Kootenay River Valley, the upper valleys of the Thompson and Fraser Rivers and the Quesnel Lake area of British Columbia contain a coniferous forest closely resembling that of the Coast Region. Western red cedar and western hemlock are the characteristic species in this interior "wet belt". Associated trees are the blue Douglas fir which is of general distribution and, in the southern parts, western white pine, western larch, grand fir and western yew. Engelmann spruce from the Subalpine Region is important in the upper Fraser Valley and is found to some extent at the upper levels of the forest in the remainder of the Region. At lower elevations in the west and in parts of the Kootenay Valley the forest grades into the Montane Region and, in a few places, into prairie grasslands.

*Deciduous Forest Region.*—A small portion of the deciduous forest, widespread in the eastern United States, occurs in southwestern Ontario between Lakes Huron, Erie and Ontario. Here, with the broadleaved trees common to the Great Lakes-St. Lawrence Region, such as sugar maple, beech, white elm, basswood, red ash, white oak and butternut, are scattered a number of other broadleaved species which have their northern limits in this locality. Among these are the tulip-tree, cucumber-tree, papaw, red mulberry, Kentucky coffee-tree, redbud, black gum, blue ash, sassafras, mockernut and pignut hickories, and scarlet, black and pin oaks. In addition, black walnut, sycamore and swamp white oak are confined largely to this Region. Conifers are few and there is only a scattered distribution of white pine, tamarack, red juniper and hemlock.

*Forest Land.*—The forest area of Canada is estimated at 1,710,788 sq. miles, about 57 p.c. of which is "productive" in the sense that it is capable of producing merchantable timber; the remainder is incapable of producing merchantable timber because of adverse climatic, soil or moisture conditions or is reserve forest land for which no inventories are available. Table 1 shows the areas of productive and non-productive forest land in each province and territory; forest land in each province classified by type of growth is given in Chapter X at p. 441.