4.40 Percentage change in death rate for each age group, by sex, 1951-72

Age group	Male	Female	Age group	Male	Female
Under 1 year	-55.5	-55.9	50-54 years	-10.6	-24.7
1- 4 years	-52.4	-55.6	55-59 "	-9.9	-27.6
5- 9 "	-50.0	-57.1	60-64 "	-6.2	-29.9
10-14 "	-37.5	-40.0	65-69 "	+1.0	-29.0
15-19 "	+1.1	-33.3	70-74 "	-2.0	-30.0
20-24 "	+1.1	-50.0	75-79 "	-9.9	-34.2
25-29 "	-16.7	-36.4	80-84 "	-9.5	-33.2
30-34 "	-19.0	-40.0	85 years and over	-9.0	-25.3
35-39 "	-12.0	-30.0			
40-44 "	-7.7	-30 0	All ages	-13.9	-20.6
45-49 "	-12.6	-31.2			

4.41 Numerical and percentage distribution of deaths by age group and sex, 1961, 1966 and 1971

Age group	Distribution									% change in death rate 1961-71
	1961			1966			1971			
	No.	%	*Rate*[1]	*No.*	%	*Rate*[1]	*No.*	%	*Rate*[1]	
Male										
Under 1 year	7,447	9.0	30.5	5,138	5.8	25.8	3,712	4.1	19.9	-34.8
1- 4 years	1,154	1.4	1.3	988	1.1	1.1	679	0.7	0.9	-30.8
5- 9 "	672	0.8	0.6	669	0.8	0.6	641	0.7	0.6	–
10-14 "	527	0.6	0.6	620	0.7	0.6	589	0.6	0.5	-16.7
15-19 "	840	1.0	1.2	1,212	1.4	1.3	1,489	1.6	1.4	+16.7
20-24 "	969	1.2	1.7	1,324	1.5	1.8	1,697	1.9	1.8	+5.9
25-29 "	895	1.1	1.5	980	1.1	1.6	1,176	1.3	1.5	–
30-34 "	1,041	1.3	1.6	1,054	1.2	1.7	1,090	1.2	1.6	–
35-39 "	1,422	1.7	2.3	1,456	1.7	2.2	1,416	1.5	2.2	-4.3
40-44 "	1,916	2.3	3.4	2,146	2.4	3.4	2,310	2.5	3.6	+5.9
45-49 "	2,993	3.6	5.8	3,111	3.5	5.7	3,523	3.8	5.7	-1.7
50-54 "	4,242	5.1	9.6	4,855	5.5	9.7	4,839	5.3	9.3	-3.1
55-59 "	5,494	6.6	15.2	6,352	7.2	15.4	6,887	7.5	14.6	-3.9
60-64 "	7,028	8.5	24.0	7,911	9.0	24.0	8,755	9.5	22.9	-4.6
65-69 "	8,545	10.3	35.7	9,226	10.5	36.2	10,279	11.2	34.7	-2.8
70-74 "	10,582	12.8	54.0	10,549	12.0	53.1	10,663	11.6	51.9	-3.9
75-79 "	10,970	13.3	81.8	11,102	12.6	79.9	11,058	12.1	79.0	-3.4
80-84 "	8,635	10.4	125.1	10,006	11.4	124.0	10,182	11.1	118.8	-5.0
85 years and over	7,337	8.9	208.9	9,214	10.5	213.4	10,838	11.8	198.5	-5.0
Total, all ages	82,709	100.0	9.0	87,913	100.0	8.7	91,823	100.0	8.5	-5.6
Female										
Under 1 year	5,493	9.4	23.7	3,822	6.2	20.2	2,644	4.0	15.1	-36.3
1-4 years	844	1.4	1.0	775	1.3	0.9	551	0.8	0.8	-20.0
5- 9 "	405	0.7	0.4	480	0.8	0.4	424	0.7	0.4	–
10-14 "	278	0.5	0.3	318	0.5	0.3	365	0.6	0.3	–
15-19 "	322	0.6	0.5	467	0.8	0.5	579	0.9	0.6	+20.0
20-24 "	342	0.6	0.6	403	0.7	0.5	559	0.9	0.6	–
25-29 "	418	0.7	0.7	384	0.6	0.6	485	0.7	0.6	-14.3
30-34 "	562	1.0	0.9	564	0.9	0.9	565	0.9	0.9	–
35-39 "	880	1.5	1.4	845	1.4	1.3	815	1.2	1.3	-7.1
40-44 "	1,099	1.9	2.0	1,293	2.1	2.0	1,290	2.0	2.1	+5.0
45-49 "	1,617	2.8	3.2	1,823	2.9	3.3	1,901	2.9	3.0	-6.2
50-54 "	2,237	3.8	5.3	2,434	3.9	5.0	2,480	3.8	4.6	-13.2
55-59 "	2,749	4.7	8.0	3,115	5.0	7.7	3,477	5.3	7.2	-10.0
60-64 "	3,725	6.4	12.8	4,064	6.6	12.2	4,345	6.6	11.0	-14.1
65-69 "	5,304	9.1	21.4	5,393	8.7	19.5	5,614	8.6	17.3	-19.2
70-74 "	7,058	12.1	34.2	7,063	11.4	30.9	7,138	10.9	28.3	-17.3
75-79 "	8,290	14.2	59.2	8,695	14.0	53.9	8,930	13.6	48.1	-18.7
80-84 "	7,871	13.5	101.2	9,048	14.6	93.6	9,763	14.9	82.4	-18.6
85 years and over	8,782	15.1	192.2	10,964	17.7	183.4	13,524	20.7	163.3	-15.0
Total, all ages	58,276	100.0	6.5	61,950	100.0	6.2	65,449	100.0	6.1	-6.2

	1961		1966		1971	
	Male	*Female*	*Male*	*Female*	*Male*	*Female*
Average age at death	59.7	63.1	62.0	65.9	63.3	68.2
Median age at death[2]	67.9	72.2	68.4	73.5	68.5	74.7

[1]Per 1,000 population per age group.
[2]The age above and below which half of the total number of annual deaths occurred.

4.42 Deaths and rate per 100,000 population according to the International Abbreviated List of 50 Causes, 1972

Abbreviated "B" List No.	Detailed List No.	Cause (eighth revision)	Deaths	Rate per 100,000 population
1	000	Cholera	–	–
2	001	Typhoid fever	–	–
3	004,006	Bacillary dysentery and amœbiasis	8	[1]
4	008,009	Enteritis and other diarrhœal diseases	298	1.4
5	010-012	Tuberculosis of respiratory system	292	1.3
6	013-019	Other tuberculosis, including late effects	161	0.7
7	020	Plague	–	–