

5.14 Reported cases of selected notifiable diseases and rate per 100,000 population, by province, 1972 (concluded)

| International List No. | Disease | Nfld. | PEI | NS | NB | Que. | Ont. | Man. | Sask. | Alta. | BC | YT | NWT | Canada |
|------------------------------------|---|-------|--------------|-------|------|------|--------------|-------|-------|-------|-------|---------|---------|--------|
| <i>Rate per 100,000 population</i> | | | | | | | | | | | | | | |
| 009.1 | Diarrhoea of the newborn, epidemic | — | 22.1 | — | — | 0.4 | ¹ | 0.1 | — | 0.3 | — | — | 5.6 | 0.4 |
| 032 | Diphtheria | — | — | — | — | — | -- | 0.5 | — | 2.9 | 0.5 | — | 2.8 | 0.3 |
| 004 | Dysentery, bacillary | 28.4 | — | — | 0.9 | 1.9 | 4.0 | 13.5 | 14.6 | 8.5 | 9.0 | — | 258.3 | 5.9 |
| 062.1 | Encephalitis, western equine | — | ¹ | — | — | — | ¹ | 0.1 | — | — | 0.3 | — | — | 0.1 |
| | Food poisoning, bacterial | — | 0.9 | — | — | 0.3 | — | 5.3 | 27.1 | 0.2 | 0.9 | — | — | 1.6 |
| 005.0 | Staphylococcal | — | 0.9 | — | — | 0.3 | ¹ | 5.3 | 27.1 | 0.2 | 0.7 | — | — | 2.4 |
| 005.1 | Botulism | — | — | — | — | -- | — | — | — | — | 0.2 | — | — | -- |
| | Hepatitis, infectious (including serum hepatitis) | 62.0 | 442.5 | 27.8 | 22.4 | 2.4 | 25.2 | 83.7 | 55.3 | 66.4 | 85.4 | 100.5 | 338.9 | 35.8 |
| 070 | Hepatitis, infectious | 62.0 | 442.5 | 27.3 | 22.4 | 1.8 | 21.6 | 79.5 | 54.2 | 62.9 | 84.3 | 89.9 | 338.9 | 33.6 |
| 999.2 | Hepatitis, serum | — | ¹ | 0.5 | — | 0.6 | 3.7 | 4.2 | 1.1 | 3.6 | 1.2 | 10.6 | — | 2.1 |
| 055 | Measles | 76.3 | — | 26.9 | 0.2 | 20.2 | 11.5 | 6.4 | 6.8 | 8.8 | 4.3 | 10.6 | 61.1 | 14.4 |
| | Meningitis, aseptic, due to enteroviruses | — | 5.3 | 0.9 | 0.5 | 1.5 | ¹ | 5.5 | 0.4 | — | 1.0 | 5.3 | 16.7 | 1.4 |
| 045.0 | Coxsackie virus | — | 0.9 | 0.3 | 0.2 | 0.6 | ¹ | 0.1 | — | — | — | — | — | 0.3 |
| 045.1 | ECHO virus | — | ¹ | — | — | 0.1 | ¹ | 0.1 | — | — | — | — | — | -- |
| 045.9 | Not specified | — | 4.4 | 0.6 | 0.3 | 0.9 | ¹ | 5.3 | 0.4 | — | 1.0 | 5.3 | 16.7 | 1.1 |
| 036 | Meningococcal infections | 8.5 | 1.8 | 1.6 | 1.2 | 1.1 | 1.2 | 6.4 | 1.9 | 1.0 | 1.5 | — | 11.1 | 1.7 |
| 056 | Rubella (German measles) | 15.4 | ¹ | 8.4 | — | 12.0 | 8.6 | 11.6 | 8.7 | 57.3 | 3.7 | 15.9 | 83.3 | 12.9 |
| | Salmonella infections, other | 11.3 | 17.7 | 8.6 | 7.0 | 16.1 | 14.6 | 11.3 | 28.8 | 26.4 | 18.6 | 10.6 | 25.0 | 16.3 |
| 003.0 | With food as vehicle | — | 14.2 | 4.8 | 1.7 | 10.9 | ¹ | 0.3 | 0.5 | 3.7 | — | — | — | 5.7 |
| 003.9 | Without mention of food as vehicle | 11.3 | 3.5 | 3.8 | 5.3 | 5.2 | 14.6 | 11.0 | 28.3 | 22.6 | 18.6 | 10.6 | 25.0 | 12.6 |
| 034 | Streptococcal sore throat and scarlet fever | 17.9 | 2,536.3 | 220.1 | 1.9 | 8.1 | 15.3 | 61.6 | 74.3 | 233.1 | 20.2 | 31.7 | 575.0 | 56.0 |
| 010,011, 012-019 | Tuberculosis | 35.2 | 4.4 | 15.4 | 12.6 | 19.8 | 13.4 | 20.5 | 18.5 | 20.7 | 22.3 | 5.3 | 136.1 | 17.9 |
| | Typhoid and paratyphoid fever | 0.2 | — | 0.1 | 0.9 | 0.7 | 0.9 | 1.4 | — | 0.1 | 0.6 | — | 16.7 | 0.7 |
| 001 | Typhoid | 0.2 | — | — | 0.9 | 0.4 | 0.5 | 1.2 | — | 0.1 | 0.4 | — | 16.7 | 0.5 |
| 002 | Paratyphoid | — | — | 0.1 | — | 0.3 | 0.4 | 0.2 | — | — | 0.2 | — | — | 0.2 |
| | Venereal diseases | 64.5 | 41.6 | 123.0 | 62.1 | 72.8 | 192.6 | 367.7 | 362.9 | 361.9 | 369.8 | 1,449.7 | 4,830.6 | 204.0 |
| 098 | Gonococcal infections | 64.3 | 37.2 | 108.9 | 59.3 | 65.3 | 172.0 | 349.7 | 344.9 | 353.2 | 353.3 | 1,444.4 | 4,794.4 | 189.9 |
| 090-097 | Syphilis | 0.2 | 4.4 | 14.1 | 2.8 | 7.4 | 20.5 | 18.1 | 17.9 | 8.6 | 16.5 | 5.3 | 36.1 | 14.0 |
| 099.0,099.1, 099.2 | Other | — | ¹ | — | — | -- | -- | — | 0.1 | — | -- | — | — | -- |
| 033 | Whooping cough | 4.5 | 45.1 | 3.3 | 8.6 | 7.5 | 4.4 | 8.6 | 5.7 | 6.3 | 4.5 | — | 2.8 | 5.9 |

¹Not reportable.