

5.24 Reported cases of selected notifiable diseases and rate per 100,000 population, by province, 1975 and 1976 (concluded)

| Year and International List No. | Disease | Nfld. | PEI | NS | NB | Que. | Ont. | Man. | Sask. | Alta. | BC | YT | NWT | Canada | |
|---------------------------------|---|------------------------------------|---------|-------|------|------|-------|-------|-------|-------|-------|---------|---------|--------|--|
| 1976 (concluded) | | <i>Rate per 100,000 population</i> | | | | | | | | | | | | | |
| 009.1 | Diarrhoea of the newborn, epidemic | — | 3.5 | — | — | — | — | — | — | — | — | — | 2.4 | 0.1 | |
| 032 | Diphtheria | 0.2 | — | — | — | — | 0.1 | 0.9 | — | 3.2 | 0.4 | — | 43.6 | 0.5 | |
| 004 | Dysentery, bacillary | 11.3 | 0.9 | — | 0.6 | 1.4 | 3.2 | 19.4 | 16.9 | 28.3 | 4.9 | 238.1 | 544.8 | 7.3 | |
| 062.1 | Encephalitis, western equine | — | — | — | — | — | — | — | — | — | — | — | — | — | |
| | Food poisoning, bacterial | 9.3 | 23.7 | 0.4 | — | 2.0 | — | 0.1 | 0.1 | 0.1 | — | — | — | 0.9 | |
| 005.0 | Staphylococcal | 9.1 | 23.7 | 0.4 | — | 1.8 | — | 0.1 | 0.1 | 0.1 | — | — | — | 1.3 | |
| 005.1 | Botulism | 0.2 | — | — | — | 0.2 | — | — | — | — | — | — | — | 0.1 | |
| | Hepatitis, infectious (including serum hepatitis) | 43.2 | 74.5 | 2.8 | 1.5 | 5.0 | 11.4 | 73.8 | 31.7 | 33.2 | 30.6 | 33.3 | 457.6 | 18.3 | |
| 070 | Hepatitis, infectious | 38.5 | 74.5 | 2.5 | 1.5 | 3.1 | 7.4 | 68.6 | 29.9 | 30.5 | 30.1 | 28.6 | 431.0 | 15.6 | |
| 999.2 | Hepatitis, serum | 47 | — | 0.3 | — | 1.9 | 4.0 | 5.2 | 1.8 | 2.7 | 0.5 | 4.7 | 26.6 | 2.7 | |
| 055 | Measles | 87.0 | — | 130.2 | 2.1 | 66.7 | 22.7 | 38.3 | 21.7 | 47.2 | 7.3 | 57.1 | 101.7 | 40.4 | |
| | Meningitis, aseptic, due to enteroviruses | 1.1 | 1.8 | 1.0 | 0.5 | 2.5 | 0.1 | 2.9 | 1.0 | 0.4 | 0.4 | 4.8 | 380.1 | 1.8 | |
| 045.0 | Coxsackie virus | — | 0.9 | — | — | 0.4 | — | 0.4 | — | — | — | — | — | 0.1 | |
| 045.1 | ECHO virus | — | — | — | — | 0.6 | — | 0.3 | 0.1 | 0.1 | — | — | — | 0.2 | |
| 045.9 | Not specified | 1.1 | 0.9 | 1.0 | 0.5 | 1.5 | 0.1 | 2.2 | 0.9 | 0.3 | 0.4 | 4.8 | 380.1 | 1.4 | |
| 036 | Meningococcal infections | 3.5 | 5.3 | 1.5 | 0.8 | 0.5 | 1.2 | 1.2 | 1.6 | 0.7 | 1.9 | 14.3 | 9.7 | 1.1 | |
| 056 | Rubella (German measles) | 3.3 | — | 5.0 | 1.5 | 14.8 | 24.7 | 16.2 | 12.4 | 40.5 | 2.8 | 9.5 | 179.2 | 18.4 | |
| | Salmonella infections, other | 13.6 | 2.6 | 9.9 | 4.4 | 8.7 | 16.5 | 8.2 | 12.8 | 17.7 | 13.0 | — | 19.4 | 12.9 | |
| 003.0 | With food as vehicle | — | 0.9 | 3.3 | 0.5 | 1.7 | — | 0.3 | — | 7.6 | — | — | — | 1.9 | |
| 003.9 | Without mention of food as vehicle | 13.6 | 1.8 | 6.6 | 4.0 | 7.0 | 16.5 | 7.9 | 12.8 | 10.1 | 13.0 | — | 19.4 | 11.7 | |
| 034 | Streptococcal sore throat and scarlet fever | 45.6 | 3,765.1 | 281.2 | 0.5 | 7.2 | 24.2 | 212.7 | 38.3 | 281.1 | 28.2 | 423.8 | 2,566.6 | 81.3 | |
| 010-018 | Tuberculosis | 20.7 | 14.0 | 9.2 | 9.7 | 12.4 | 8.2 | 18.3 | 15.5 | 11.4 | 12.9 | 38.1 | 106.5 | 11.4 | |
| | Typhoid and paratyphoid fever | 0.4 | — | 0.9 | 1.8 | 2.8 | 0.6 | 0.2 | 0.6 | 0.1 | 0.3 | — | — | 1.1 | |
| 001 | Typhoid | 0.4 | — | — | 1.1 | 2.7 | 0.5 | 0.2 | — | 0.1 | 0.1 | — | — | 1.0 | |
| 002 | Paratyphoid | — | — | 0.9 | 0.8 | 0.1 | 0.1 | — | 0.6 | 0.1 | 0.2 | — | — | 0.2 | |
| | Venereal diseases | 144.1 | 145.5 | 117.2 | 46.3 | 78.3 | 241.5 | 492.2 | 401.5 | 496.5 | 407.7 | 1,761.9 | 3,770.0 | 247.4 | |
| 098 | Gonococcal infections | 141.9 | 142.0 | 112.3 | 44.6 | 67.0 | 209.5 | 475.0 | 394.4 | 491.5 | 394.5 | 1,761.9 | 3,760.3 | 229.5 | |
| 090-097 | Syphilis | 2.2 | 3.5 | 4.9 | 1.7 | 11.3 | 31.8 | 6.6 | 6.4 | 5.0 | 13.2 | — | 9.7 | 17.4 | |
| 099.0,099.1, | | | | | | | | | | | | | | | |
| 099.2 | Other | — | — | — | — | — | 0.2 | 10.5 | 0.7 | — | — | — | — | 0.6 | |
| 033 | Whooping cough | 38.3 | 209.5 | 17.5 | 2.6 | 4.6 | 17.9 | 9.2 | 11.6 | 19.6 | 3.1 | 4.8 | 2.4 | 13.2 | |

¹Not reportable.

²Tuberculosis figures are provisional.

³Excludes 11 cases of syphilis, type undetermined.

⁴Excludes 22 cases of syphilis, type undetermined.