

PROPORTION OF DEATHS FROM TYPHOID FEVER PER 1,000 OF  
POPULATION IN THE FOLLOWING PLACES IN 1890.

CITIES AND TOWNS.	Per 1,000.	CITIES AND TOWNS.	Per 1,000.
Winnipeg .....	1·09	Montreal.....	0·35
St. Johns, Que .....	1·05	Woodstock, Ont. ....	0·35
Brantford.....	0·86	Quebec.....	0·32
St. Thomas.....	0·67	Belleville .....	0·30
Toronto .....	0·65	Windsor.....	0·29
Kingston.....	0·52	Guelph.....	0·28
Hamilton.....	0·47	St. John, N.B.....	0·26
Fredericton.....	0·46	Charlottetown.....	0·26
Chatham, Ont.....	0·44	Peterborough.....	0·21
Victoria, B.C.....	0·44	London.....	0·19
St. Hyacinthe.....	0·43	Halifax .....	0·18
Ottawa.....	0·43	Galt .....	0·13
Three Rivers.....	0·36	Belleville.....	0·11
Hull.....	0·36		

There was an increase in the rate per 1,000 of the aggregate population of all the places making returns from ·41 to ·43, but if Toronto is taken out the rate is reduced to ·37. Even this is higher than it ought to be; in England and Wales in 1889 it was only ·17. The proportion in Ontario in 1889 was ·26 per 1,000 inhabitants.

148. The following table shows how the assigned causes of death, in the places making returns, were distributed among the several classes. The classification is based upon that established by the Registrar-General of England :—

Causes of  
death by  
classes.

CITIES AND TOWNS.	Total Deaths.	Causes of death by classes.								
		Zymotic.	Parasitic.	Dietetic.	Constitutional.	Developmental.	Local.	Violence.	Ill-defined and not specified causes.	
Montreal.....	6,209	1,296	1	12	888	352	2,636	156	868	
Toronto.....	2,919	515	9	4	459	211	1,302	92	327	
Quebec.....	2,116	555	4	4	256	235	783	34	249	
Hamilton .....	867	125	4	141	65	425	31	76		
Ottawa.....	948	217	6	148	91	380	33	73		
St. John, N.B.....	741	126	130	64	320	37	64			
Halifax.....	1,089	306	1	2	155	67	448	21	89	
London.....	359	57	67	43	162	8	22			
Winnipeg.....	403	117	2	1	63	15	154	21	30	
Victoria, B.C.....	275	44	1	2	44	17	124	17	26	
Kingston.....	383	58	1	55	60	144	18	47		
Hull.....	384	64	22	17	111	8	162			