Causes of Death.—About nine-tenths of deaths in Canada are due to the 28 causes specified in Table 25. About three-quarters are due to the 10 leading causes: diseases of the heart, cancer, intracranial lesions, violent deaths, nephritis, diseases of early infancy, pneumonia, tuberculosis, influenza and diseases of the arteries.

The classification of the causes of death is according to the revision of the International List of 1938, which was first used in Canada in 1941. Each revision of the International List makes continuity of classification difficult. This applies especially to diseases of the heart, intracranial lesions (cerebral hæmorrhage) and diseases of the arteries.

The rise in the average age at death has been noted above. Causes of death that affect mainly children and young adults have declined. Diphtheria, for example, has very nearly been wiped out. Tuberculosis has also been greatly reduced. On the other hand, the ageing of the population increases the proportion of deaths from the causes that affect mainly older people. Thus, cancer, nephritis and diseases of the heart now account for a substantially larger proportion of all deaths, than they used to.

25.—Deaths and Death Rates per 100,000 Population, by Principal Causes, 1941-45 Note.—Figures for 1944 and 1945 are by residence; for previous years by place of occurrence.

Inter- national List No.1	Cause of Death	1941	1942	1943	1944	1945
		NUMBERS OF DEATHS				
$\begin{array}{c} 1,\ 2\\ 8\\ 9\\ 10\\ 13\\ 35\\ 45-55\\ 61\\ 73\\ 83\\ 86\\ 90-95\\ 96,\ 97,\\ 99,\ 102\\ 106\\ 107-109\\ 119,\ 120\\ 122\\ 130-132\\ 137\\ 140-150\\ 158-161\\ 162\\ 163,\ 164\\ 166-198\\ \end{array}$	Typhoid fever, including paratyphoid. Scarlet fever. Whooping cough. Diphtheria. Tuberculosis, respiratory system. Tuberculosis, other organs. Influenza. Measles. Cancer and other malignant tumors. Diabetes mellitus. Anæmias. Intracranial lesions of vascular origin. Convulsions (under 5 years of age). Diseases of the heart. Diseases of the arteries. Bronchitis. Pneumonia. Diarrhœa and enteritis. Appendicitis. Hernia, intestinal obstruction Nephritis. Diseases of the prostate. Puerperal causes. Congenital malformations. Diseases peculiar to first year of life. Senility. Suicides. Violent deaths (suicides excepted). Other specified causes.	165 117 437 240 5,002 1,070 2,411 3,417 2,140 408 9,034 199 26,602 2,266 394 5,955 2,319 908 7,399 901 1,051 908 7,399 901 6,502 1,970 1,0	108 1290 560 256 4,947 1,033 1,227 13,654 2,242 2,242 27,529 2,702 383 5,778 2,400 824 912 7,233 855 818 6,029 1,650 839 7,332 11,493	758 7,516 11,289	131 115 337 309 4,705 1,019 1,864 239 14,271 2,362 25 9,089 15 29,148 2,349 431 5,940 2,695 9,117 7,124 6,655 1,69	10 71 476 476 98 1,08 1,08 9,42 14,43 2,41 35 9,42 2,01 2,01 86 67 86 66,92 84 66,92 1,62 7,70 1,62 7,70 1,62 7,70 1,62 1,63 1,63 1,63 1,63 1,63 1,63 1,63 1,63
199, 200	Totals, Specified Causes	113,601 1,038 114,639	973	884	115,243 809 116,052	87

For footnote, see end of table, p. 167.